arXiv:1603.01832v1 [math.CA] 6 Mar 2016

DUNKL HARMONIC ANALYSIS AND FUNDAMENTAL SETS OF FUNCTIONS ON THE UNIT SPHERE

Roman A. Veprintsev

Abstract. Using Dunkl theory, we introduce into consideration some weighted L_p-spaces on $[-1,1]$ and on the unit Euclidean sphere $\mathbb{S}^{d-1}$, $d \geq 2$. Then we define a family of linear bounded operators $\{V_\kappa^p(x)\colon x \in \mathbb{S}^{d-1}\}$ acting from the L_p-space on $[-1,1]$ to the L_p-space on $\mathbb{S}^{d-1}$, $1 \leq p < \infty$. We establish a necessary and sufficient condition for a function g belonging to the L_p-space on $[-1,1]$ such that the family of functions $\{V_\kappa^p(x;g)\colon x \in \mathbb{S}^{d-1}\}$ is fundamental in the L_p-space on $\mathbb{S}^{d-1}$.

Key words and phrases: fundamental set, unit sphere, Dunkl theory, Dunkl intertwining operator, Funk – Hecke formula for κ-spherical harmonics

MSC 2010: 42B35, 42C05, 42C10

1. Introduction and preliminaries

In this section we introduce some basic definitions and notions of general Dunkl theory (see, e.g., [2–6]); for a background on reflection groups and root systems the reader is referred to [6, 8].

Let $\mathbb{N}_0$ be the set of nonnegative integers, let $\mathbb{R}^d$ be the d-dimensional real Euclidean space of all d-tuples of real numbers. For $x \in \mathbb{R}^d$, we write $x = (x_1, \ldots, x_d)$. The inner product of x, $y \in \mathbb{R}^d$ is denoted by $\langle x, y\rangle = \sum\limits_{i=1}^{d} x_i y_i$, and the norm of x is denoted by $\|x\| = \sqrt{\langle x, x\rangle}$. Let $\mathbb{S}^{d-1} = \{x\colon \|x\| = 1\}$ be the unit sphere in $\mathbb{R}^d$, $d \geq 2$. Denote by $d\omega$ the usual Lebesgue measure on $\mathbb{S}^{d-1}$.

For a nonzero vector $v \in \mathbb{R}^d$, define the reflection s_v by

$$s_v(x) = x - 2\frac{\langle x, v\rangle}{\|v\|^2}\, v, \quad x \in \mathbb{R}^d.$$

Each reflection s_v is contained in the orthogonal group $O(\mathbb{R}^d)$.

Recall that a finite set $R \subset \mathbb{R}^d \setminus \{0\}$ is called a root system if the following conditions are satisfied:

(1) $R \cap \mathbb{R}v = \{\pm v\}$ for all $v \in R$; (2) $s_v(R) = R$ for all $v \in R$.

The subgroup $G = G(R) \subset O(\mathbb{R}^d)$ which is generated by the reflections $\{s_v\colon v \in R\}$ is called the reflection group associated with R. It is known that the reflection group G is finite and the set of reflections contained in G is exactly $\{s_v\colon v \in R\}$.

Each root system R can be written as a disjoint union $R = R_+ \cup (-R_+)$, where R_+ and $-R_+$ are separated by a hyperplane through the origin. Such a set R_+ is called a positive subsystem. Its choice in not unique.

A nonnegative function κ on a root system R is called a multiplicity function if it is G-invariant, i.e. $\kappa(v) = \kappa(g(v))$ for all $v \in R$, $g \in G$.

Note that definitions given below do not depend on the special choice of R_+, thanks to the G-invariance of κ.

The Dunkl operators are defined by

$$\mathcal{D}_i f(x) = \frac{\partial f(x)}{\partial x_i} + \sum_{v \in R_+} \kappa(v) \frac{f(x) - f(s_v(x))}{\langle v, x \rangle} v_i, \quad 1 \le i \le d.$$

In case $\kappa = 0$, the Dunkl operators reduce to the corresponding partial derivatives. These operators were introduced and first studied by C. F. Dunkl.

Let

$$\lambda_\kappa = \gamma_\kappa + \frac{d-2}{2}, \qquad \gamma_\kappa = \sum_{v \in R_+} \kappa(v),$$

let w_κ denote the weight function on $\mathbb{S}^{d-1}$ defined by

$$w_\kappa(x) = \prod_{v \in R_+} |\langle v, x \rangle|^{2\kappa(v)}, \quad x \in \mathbb{S}^{d-1}.$$

The weight function w_κ is a positively homogeneous G-invariant function of degree $2\gamma_\kappa$. In case $\kappa = 0$, w_κ is identically equal to 1.

Suppose Π^d is the space of all polynomials in d variables with complex coefficients, $\mathcal{P}_n^d$ is the subspace of homogeneous polynomials of degree $n \in \mathbb{N}_0$ in d variables.

C. F. Dunkl has proved in [5] that there exists a unique linear isomorphism V_κ of Π^d such that

$$V_\kappa(\mathcal{P}_n^d) = \mathcal{P}_n^d, \ \ n \in \mathbb{N}_0, \qquad V_\kappa 1 = 1, \qquad \mathcal{D}_i V_\kappa = V_\kappa \frac{\partial}{\partial x_i}, \ \ 1 \le i \le d.$$

This operator is called the Dunkl intertwining operator. The operator V_κ was studied by many mathematicians (for example, C. F. Dunkl, M. Rösler, K. Trimèche, Y. Xu). If $\kappa = 0$, then V_κ is the identity operator.

Throughout this paper, we assume that $p \in [1, \infty)$ and $\lambda_\kappa > 0$. In particular, it follows that $\gamma_\kappa > 0$ if $d = 2$.

To explain our main result of the present paper, we need to introduce some weighted L_p-spaces and one family of linear operators.

Denote by $L_{\kappa,p}(\mathbb{S}^{d-1})$ the space of complex-valued Lebesgue measurable functions f on $\mathbb{S}^{d-1}$ with finite norm

$$\|f\|_{\kappa,p,\mathbb{S}^{d-1}} = \Big(\int_{\mathbb{S}^{d-1}} |f(x)|^p \, d\sigma_\kappa(x) \Big)^{1/p}, \qquad d\sigma_\kappa(x) = a_\kappa \, w_\kappa(x) \, d\omega(x),$$

where the normalizing constant a_κ satisfies $a_\kappa \int_{\mathbb{S}^{d-1}} w_\kappa \, d\omega = 1$. The space $L_{\kappa,2}(\mathbb{S}^{d-1})$ is a complex Hilbert space with the inner product

$$\langle f, h \rangle_{\kappa,\mathbb{S}^{d-1}} = \int_{\mathbb{S}^{d-1}} f(x) \overline{h(x)} \, d\sigma_\kappa(x).$$

We also introduce the space $L_{\kappa,\infty}(\mathbb{S}^{d-1})$ composed of all complex-valued Lebesgue measurable functions defined on $\mathbb{S}^{d-1}$ which are σ_κ-measurable and σ_κ-essentially bounded. Because w_κ is ω-a.e. nonzero on $\mathbb{S}^{d-1}$, the above notions coincides with the one of w_κ-measurable and w_κ-essentially bounded function, respectively.

Let $\lambda > 0$. Suppose $L_{p,\lambda}[-1,1]$ is the space of complex-valued Lebesgue measurable functions g on the segment $[-1,1]$ with finite norm

$$\|g\|_{\lambda,p,[-1,1]} = \Big(c_\lambda \int_{-1}^{1} |g(t)|^p \,(1-t^2)^{\lambda-1/2}\,dt\Big)^{1/p}, \qquad c_\lambda = \Big(\int_{-1}^{1} (1-t^2)^{\lambda-1/2}\,dt\Big)^{-1}.$$

The Gegenbauer polynomials C_n^λ (see, e.g., [1, p. 302]) are orthogonal with respect to the weight function $(1-t^2)^{\lambda-1/2}$. For a function $g \in L_{p,\lambda}[-1,1]$, its Gegenbauer expansion takes the form

$$g(t) \sim \sum_{n=0}^{\infty} \Lambda_{n,\lambda}(g) \frac{n+\lambda}{\lambda} C_n^\lambda(t) \quad \text{with} \quad \Lambda_{n,\lambda}(g) = \frac{c_\lambda}{C_n^\lambda(1)} \int_{-1}^{1} g(t) C_n^\lambda(t) (1-t^2)^{\lambda-1/2}\,dt, \quad (1)$$

since $\|C_n^\lambda\|^2_{\lambda,2,[-1,1]} = C_n^\lambda(1)\lambda/(n+\lambda)$.

Theorem 13.17 in [7] says that $L_{\kappa,q}(\mathbb{S}^{d-1}) \subset L_{\kappa,p}(\mathbb{S}^{d-1})$ and $L_{q,\lambda}[-1,1] \subset L_{p,\lambda}[-1,1]$ for $1 \le p < q < \infty$.

Let us now consider the operators

$$V_\kappa^p(x)\colon L_{p,\lambda_\kappa}[-1,1] \to L_{\kappa,p}(\mathbb{S}^{d-1})$$

depending on $x \in \mathbb{S}^{d-1}$ which are defined by the following rule:

$$V_\kappa^p(x; g, y) = V_\kappa\big[g(\langle x, \cdot\rangle)\big](y), \quad y \in \mathbb{S}^{d-1} \quad \big(g \in L_{p,\lambda_\kappa}[-1,1]\big).$$

They are linear bounded operators with operator norm 1. It follows from the properties of V_κ known from the general Dunkl theory (see especially the proof of Theorem 7.4.2 in [2]) that these operators are well defined.

The main result of this paper is a generalization of Theorem 2.4 in [10]. More precisely, we establish a necessary and sufficient condition for a function $g \in L_{p,\lambda_\kappa}[-1,1]$ such that the family of functions

$$\mathcal{M}_\kappa^p(g) = \{V_\kappa^p(x; g)\colon x \in \mathbb{S}^{d-1}\} \quad (2)$$

is fundamental in the space $L_{\kappa,p}(\mathbb{S}^{d-1})$. This fact is formulated and proved in section 3.

Recall that a set $\mathcal{F}$ in a Banach space $\mathcal{E}$ is said to be fundamental if the linear span of $\mathcal{F}$ is dense in $\mathcal{E}$. To prove the main result, we use a consequence of the Hahn–Banach theorem related to fundamentality of sets in normed linear spaces. We include it as a separate lemma for convenience.

Lemma 1. *Let $\mathcal{F}$ be a subset of a Banach space $\mathcal{E}$. In order that $\mathcal{F}$ be fundamental in $\mathcal{E}$, it is necessary and sufficient that $\mathcal{F}$ not be annihilated by a nonzero bounded linear functional on $\mathcal{E}$.*

2. Some facts of Dunkl harmonic analysis on the unit sphere

The Dunkl Laplacian Δ_κ is defined by

$$\Delta_\kappa = \sum_{i=1}^{d} \mathcal{D}_i^2$$

and it plays the role similar to that of the ordinary Laplacian. It reduces to the ordinary Laplacian provided that $\kappa = 0$.

A κ-harmonic polynomial P of degree $n \in \mathbb{N}_0$ in d variables is a homogeneous polynomial $P \in \mathcal{P}_n^d$ such that $\Delta_\kappa P = 0$. Its restriction to the unit sphere is called the κ-spherical harmonic of degree n in d variables. Denote by $\mathcal{A}_n^d(\kappa)$ the space of κ-spherical harmonics of degree n in d variables. The κ-spherical harmonics of different degrees turn out to be orthogonal with respect to the weighted inner product $\langle \cdot, \cdot \rangle_{\kappa, \mathbb{S}^{d-1}}$ [3, Theorem 1.6].

Let $C(\mathbb{S}^{d-1})$ be the space of complex-valued continuous functions on $\mathbb{S}^{d-1}$.

Lemma 2. *The set* $\bigcup\limits_{n=0}^{\infty} \mathcal{A}_n^d(\kappa)$ *is fundamental in* $C(\mathbb{S}^{d-1})$ *and in* $L_{\kappa,p}(\mathbb{S}^{d-1})$, $1 \le p < \infty$.

P r o o f. Theorem 3.14 in [11] states that the space $C(\mathbb{S}^{d-1})$ is dense in $L_{\kappa,p}(\mathbb{S}^{d-1})$ for $1 \le p < \infty$. So it is sufficient to show that $\bigcup\limits_{n=0}^{\infty} \mathcal{A}_n^d(\kappa)$ is fundamental in $C(\mathbb{S}^{d-1})$.

By the Weierstrass approximation theorem, if f is continuous on $\mathbb{S}^{d-1}$, then it can be uniformly approximated by polynomials restricted to $\mathbb{S}^{d-1}$. According to [3, Theorem 1.7], these restrictions belong to the linear span of $\bigcup\limits_{n=0}^{\infty} \mathcal{A}_n^d(\kappa)$. Thus, $\bigcup\limits_{n=0}^{\infty} \mathcal{A}_n^d(\kappa)$ is fundamental in $C(\mathbb{S}^{d-1})$. □

The above proof is analogous to that of Corollary 2.3 in [12].

Lemma 3. *Let* $g \in L_{p,\lambda_\kappa}[-1,1]$, $1 \le p < \infty$. *Then for every* $Y_n^\kappa \in \mathcal{A}_n^d(\kappa)$,

$$\int_{\mathbb{S}^{d-1}} V_\kappa^p(x; g, y)\, Y_n^\kappa(y)\, d\sigma_\kappa(y) = \Lambda_{n,\lambda_\kappa}(g)\, Y_n^\kappa(x), \quad x \in \mathbb{S}^{d-1}, \tag{3}$$

where the constant $\Lambda_{n,\lambda_\kappa}(g)$ *is defined from* (1).

Equality (3) is the Funk – Hecke formula for κ-spherical harmonics written in our setting and designations (cf. [2, Theorem 7.2.7], [13, Theorem 2.1]).

3. Main result: proof and its consequence

We can now state and prove the main theorem.

Theorem 1. *Let* $d \ge 2$, $1 \le p < \infty$. *Fix a root system* R *in* $\mathbb{R}^d$ *and a multiplicity function* κ *on* R. *Let* $g \in L_{p,\lambda_\kappa}[-1,1]$. *In order that the set* $\mathcal{M}_\kappa^p(g)$ (2) *be fundamental in* $L_{\kappa,p}(\mathbb{S}^{d-1})$, *it is necessary and sufficient that* $\Lambda_{n,\lambda_\kappa}(g) \ne 0$ (1) *for every* $n \in \mathbb{N}_0$.

P r o o f. We first prove that the condition is sufficient. Let Φ be a bounded linear functional on $L_{\kappa,p}(\mathbb{S}^{d-1})$ which annihilates $\mathcal{M}_\kappa^p(g)$. According to the Riesz representation theorem [11, Theorem 6.16], Φ can be written as follows: $\Phi(\,\cdot\,) = \langle\, \cdot\,, h\rangle_{\kappa,\mathbb{S}^{d-1}}$, where $h \in$ $\in L_{\kappa,q}(\mathbb{S}^{d-1})$ and q is the exponent conjugate to p ($p^{-1} + q^{-1} = 1$; $q = \infty$ whenever $p = 1$). Then the annihilating property of Φ reduces to

$$\int_{\mathbb{S}^{d-1}} V_\kappa^p(x; g, y)\, \overline{h(y)}\, d\sigma_\kappa(y) = 0, \quad x \in \mathbb{S}^{d-1}.$$

Next, we multiply both sides of the previous equality by $Y_n^\kappa(x) \in \mathcal{A}_n^d(\kappa)$, $n \in \mathbb{N}_0$, and integrate the resulting expression with respect to the measure $d\sigma_\kappa$. Hölder's inequality implies that $V_\kappa^p(x;g,y)\,\overline{h(y)}\,Y_n^\kappa(x)$ is $\sigma_\kappa \times \sigma_\kappa$-integrable over $\mathbb{S}^{d-1} \times \mathbb{S}^{d-1}$, and hence, using the Fubini theorem to interchange the order of integration, we get

$$\int_{\mathbb{S}^{d-1}} \overline{h(y)} \left(\int_{\mathbb{S}^{d-1}} V_\kappa^p(x;g,y)\, Y_n^\kappa(x)\, d\sigma_\kappa(x) \right) d\sigma_\kappa(y) = 0.$$

Using the symmetric relation [9, formula (7)]

$$V_\kappa\big[g(\langle x,\cdot\rangle)\big](y) = V_\kappa\big[g(\langle y,\cdot\rangle)\big](x) \quad \sigma_\kappa \times \sigma_\kappa\text{-a.e. on } \mathbb{S}^{d-1}\times\mathbb{S}^{d-1}$$

and the Funk – Hecke formula (3), we obtain

$$\Lambda_{n,\lambda_\kappa}(g)\,\langle Y_n^\kappa, h\rangle_{\kappa,\mathbb{S}^{d-1}} = 0, \qquad Y_n^\kappa \in \mathcal{A}_n^d(\kappa), \quad n \in \mathbb{N}_0.$$

It follows from the condition that

$$\langle Y_n^\kappa, h\rangle_{\kappa,\mathbb{S}^{d-1}} = 0, \qquad Y_n^\kappa \in \mathcal{A}_n^d(\kappa), \quad n \in \mathbb{N}_0.$$

Thus, Φ annihilates $\bigcup\limits_{n=0}^{\infty} \mathcal{A}_n^d(\kappa)$. By continuity of Φ and Lemma 2, $\Phi = 0$ on $L_{\kappa,p}(\mathbb{S}^{d-1})$. Therefore, the set $\mathcal{M}_\kappa^p(g)$ is fundamental in $L_{\kappa,p}(\mathbb{S}^{d-1})$ by Lemma 1.

Let us now prove that the condition described in the theorem is necessary. Assume, to reach a contradiction, that there exists an index $m \in \mathbb{N}_0$ such that $\Lambda_{m,\lambda_\kappa}(g) = 0$. Select any nontrivial κ-spherical harmonic $Y_m^\kappa \in \mathcal{A}_m^d(\kappa)$ and consider a measure μ defined on the Lebesgue subsets $\mathcal{L}$ of $\mathbb{S}^{d-1}$ by the rule

$$\mu(B) = \int_B Y_m^\kappa(x)\,\sigma_\kappa(x), \quad B \in \mathcal{L}.$$

This measure is nontrivial by its definition.

Using the Funk – Hecke formula (3), we obtain

$$\begin{aligned}\int_{\mathbb{S}^{d-1}} V_\kappa^p(x;g,y)\, d\mu(y) &= \int_{\mathbb{S}^{d-1}} V_\kappa^p(x;g,y)\, Y_m^\kappa(y)\, d\sigma_\kappa(y) \\ &= \Lambda_{m,\lambda_\kappa}(g)\, Y_m^\kappa(x) = 0, \quad x \in \mathbb{S}^{d-1}.\end{aligned}$$

Thus, the nontrivial bounded linear functional Φ_1 on $L_{\kappa,p}(\mathbb{S}^{d-1})$ given by $\Phi_1(f) = \int_{\mathbb{S}^{d-1}} f\, d\mu$ annihilates $\mathcal{M}_\kappa^p(g)$. By Lemma 1, $\mathcal{M}_\kappa^p(g)$ is not fundamental in $L_{\kappa,p}(\mathbb{S}^{d-1})$. This contradicts our assumption. □

The above proof is exactly like that of Theorem 2.4 in [10]. Using the scheme of the proof of the theorem, one can prove the following result.

Corollary 1. *Let $d \geq 2$, $s \geq 1$, $1 \leq p < \infty$. Fix a root system R in $\mathbb{R}^d$ and a multiplicity function κ on R. Let $g_1, \ldots, g_s \in L_{p,\lambda_\kappa}[-1,1]$. In order that the set $\bigcup\limits_{i=1}^{s} \mathcal{M}_\kappa^p(g_i)$ be fundamental in $L_{\kappa,p}(\mathbb{S}^{d-1})$, it is necessary and sufficient that $\sum\limits_{i=1}^{s} |\Lambda_{n,\lambda_\kappa}(g_i)| \neq 0$ for every $n \in \mathbb{N}_0$.*

REFERENCES

1. Andrews, G.E., Askey, R. and Roy, R., *Special functions*. Cambridge University Press, 1999.
2. Dai, F. and Xu, Y., *Approximation theory and harmonic analysis on spheres and balls*. Springer, Berlin–New York, 2013.
3. Dunkl, C.F., Reflection groups and orthogonal polynomials on the sphere. *Math. Z.* 197: 33–60, 1988.
4. Dunkl, C.F., Differential-difference operators associated to reflection groups. *Trans. Amer. Math. Soc.* 311(1): 167–183, 1989.
5. Dunkl, C.F., Integral kernels with reflection group invariance. *Can. J. Math.* 43(6): 1213–1227, 1991.
6. Dunkl, C.F. and Xu, Y., *Orthogonal polynomials of several variables*. 2nd ed., Cambridge University Press, 2014.
7. Hewitt, E. and Stromberg, K., *Real and abstract analysis*. Springer-Verlag, New York, 1965.
8. Humphreys, J.E., *Reflection groups and Coxeter groups*. Cambridge University Press, 1990.
9. Li, Zh., Song, F., Inversion formulas for the spherical Radon–Dunkl transform. *SIGMA*. 5: 025, 15 pages, 2009.
10. Menegatto, V.A., Fundamental sets of functions on spheres. *Methods Appl. Anal.* 5(4): 387–398, 1998.
11. Rudin, W., *Real and complex analysis*. 3rd ed., McGraw-Hill, New York, 1987.
12. Stein, E.M. and Weiss, G., *Introduction to Fourier analysis on Euclidean spaces*. Princeton University Press, Princeton, 1971.
13. Xu, Y., Funk – Hecke formula for orthogonal polynomials on spheres and on balls. *Bull. London Math. Soc.* 32: 447–457, 2000.

Independent researcher, Uzlovaya, Russia
E-mail address: **veprintsevroma@gmail.com**